

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/353752310>

Study on the utilization, chemical composition, and insecticidal activity of nutmeg essential oil (*Myristica fragnans* Houtt) against fruit flies, *Bactrocera* spp. (Diptera: Tephrit...

Article in *Ecology, Environment and Conservation* · August 2021

CITATIONS

0

READS

127

5 authors, including:



I Wayan Susila

Udayana University

50 PUBLICATIONS 48 CITATIONS

[SEE PROFILE](#)



I Wayan Supartha

Udayana University

83 PUBLICATIONS 98 CITATIONS

[SEE PROFILE](#)



I Kadek Wisma Yudha

Udayana University

15 PUBLICATIONS 17 CITATIONS

[SEE PROFILE](#)



Putu Angga Wiradana

Universitas Dhyana Pura Bali

50 PUBLICATIONS 37 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



CURRENT STATUS OF *Liriomyza* spp. (DIPTERA: AGROMYZIDAE) AS AN INVASIVE PEST IN VEGETABLE PLANTS IN BALI [View project](#)



Invasive Pest, *Phenacoccus manihoti* and Natural Enemies Fauna Associated with Several Cassava Cultivars in Bali [View project](#)

Study on the utilization, chemical composition, and insecticidal activity of nutmeg essential oil (*Myristica fragrans* Houtt) against fruit flies, *Bactrocera* spp. (Diptera: Tephritidae)

I Wayan Susila^{1*}, I. Wayan Supartha¹, I. Ketut Sumiartha¹, I. Kadek Wisma Yudha² and Putu Angga Wiradana³

¹ Laboratory of Integrated Pest Management (IPMLab), Faculty of Agriculture, Udayana University, Denpasar City, 80231, Indonesia

² Graduate Program of Department of Dry Land Agriculture, Faculty of Agriculture, Udayana University, Denpasar City, 80231, Indonesia

³ Study Program of Biology, Faculty of Health, Science and Technology, Universitas Dhyana Pura, North Kuta, Badung Regency (80361) Bali Province, Indonesia

(Received 12 February, 2021; Accepted 21 April, 2021)

ABSTRACT

This study aims to determine the activity of the essential oil nutmeg (*Myristica fragrans* Houtt) against fruit flies *Bactrocera* spp. The attractant treatments used were 1) pure nutmeg essential oil (NEO), 2) petrogenol (800 g/l methyl eugenol) and 3) sterile water as a control. The variables observed were the type of species, sex, number of fruit flies captured, and activity period of NEO and Petrogenol in the field scale. The chemical composition of NEO was investigated by Gas Chromatography-Mass Spectrometry (GC-MS). The results of the research indicate the period of activity of NEO and Petrogenol in the field is more than 43 days with an average of fruit fly attracted was 12.22 adults/day and 17.04 adults/day respectively. There are three species of fruit flies attracted namely *Bactrocera carambolae*, *B. dorsalis*, and *B. umbrosa*. Among those species, *B. carambolae* and *B. dorsalis* were dominantly attracted. GC-MS analysis shows that the highest chemical composition of NEO is obtained by 4-allyl-1,2-dimethoxybenzene-carboxylate (Methyl eugenol) (8.33%) which has the potential to be developed as a fruit fly attractant.

Key words : Natural attractant, Essential oil, Fruit fly, Methyl eugenol, Biopesticides.

Introduction

Fruit flies (Diptera: Tephritidae) are one of the main pests that can harm the productivity of fruit and vegetable commodities in Indonesia. Loss of fruit caused by the fruit fly maggots eating the fruit pulp rots the fruit before it ripens (Radonjic *et al.*, 2019). Fruit flies can cause direct damage to fruits and vegetables which will cause loss of agricultural produc-

tion by up to 90-100% depending on the fruit fly population, locality, variety, and season (Sarwar *et al.*, 2014). In general, the loss of agricultural output due to fruit flies can vary between 30-100% (Dhillon *et al.*, 2005). About 35% of species attack soft-textured fruit, including fruit cultivated by farmers (White and Elson-Harris 1992). In India, about 50% of the Cucurbitaceae are attacked by *Bactrocera cucurbitae* (Jaleel *et al.*, 2020). Furthermore, reported

by Siwi *et al.* (2006) there are 16 species of fruit flies that are considered important pests of vegetable and fruit commodities in Indonesia.

To date, various attempts have been made to control fruit flies in Indonesia. However, the use of pesticides is still predominantly carried out on a field scale by many fruit and vegetable farmers (Ngowi *et al.*, 2007; Thanh Mai *et al.*, 2018; Wilson and Tisdell 2001). The unwise use of pesticides is considered an unsustainable effort for the environment (Aktar *et al.*, 2009); Prastika *et al.*, 2021; Afifah *et al.*, 2020), environmental hazards such as harmful side effects on beneficial insects (Sanchez-Bayo 2011), and human health (Singh *et al.*, 2018); Ibrahim *et al.*, 2020a; Ibrahim *et al.*, 2020b).

Essential oil (EO) is a potential source that can be developed as a promising bioinsecticide because the chemical content of secondary metabolites is dominated by monoterpenes, sesquiterpenes, phenylpropanoids, and hydrocarbons (Setiawan *et al.*, 2016; Butnariu and Sarac 2018; Kasman *et al.* 2020) Recently, the development of essential oils isolated from plants has great promise for use in pest control. Plants that have the potential to be used as this material are Nutmeg (*Myristica fragrans* Houtt). Since ancient times, local people in Indonesia have used *Myristica fragrans* seeds as a food spice, flavor, and traditional medicine for headaches and fevers (Muchtari *et al.*, 2010). Nutmeg essential oil also has antimicrobial, antiseptic, antiparasitic, anti-inflammatory, and antioxidant properties (Muchtari *et al.*, 2010; Baser and Buchbauer 2015). The report also shows that the essential oil concentration in nutmeg ranges from 5 - 15% and its secondary metabolite components are dominated by terpene hydrocarbons, oxygenated terpenes, and aromatic ethers (Gupta *et al.*, 2013; Lanari *et al.*, 2018; Baser and Buchbauer 2015).

The use of attractants or natural essential oils (EO) is one component of an integrated pest management system that is environmentally friendly and sustainable (Damos *et al.*, 2015). This method is useful as an effective, efficient, and environmentally friendly way to control fruit flies in the field. Until now, this type of synthetic attractant has been widely sold in Indonesia under the brand name Petrogenol 800 g/L methyl eugenol. Apart from synthetic attractants, there are also attractants sourced from *Ocimum* spp. plant. According to Kardinan *et al.* (2020) controlling fruit flies using *Ocimum* spp. as an attractant can increase produc-

tion by up to 73%.

Based on this, this study was conducted to determine the activity of the essential oil nutmeg (*Myristica fragrans* Houtt) against fruit flies (*Bactrocera dorsalis* Complex). This research was conducted to develop natural attractant materials that have the potential to be developed in the biological control of the agricultural sector.

Materials and Methods

Study area

The experiment was conducted in the Laboratory of Integrated Pest Management, Faculty of Agriculture, Udayana University and field the experiment was conducted in Denpasar City, Bali, Indonesia from May to September 2020.

Extraction of Essential Oils (EOs)

The nutmeg oil extract was made using the Steam Distillation method. In short, the nutmeg which consists of seeds, mace, and pulp is uniformly cut and dried for 5 - 7 days. After drying, the nutmegs are put into a kettle that has been filled with water as high as 5 cm. The kettle is tightly closed and heated. The heating process lasts for 24 hours. The essential oil and water produced are separated using a separating funnel. The essential oil produced in this distillation process is used to test (bioassay) and analyze bioactive materials by GC-MS.

GC-MS Analysis

The bioactive ingredients contained in essential oil extracts were analyzed using Gas chromatography-mass spectrometry (GC-MS) based on company protocol (Bernardi *et al.*, 2016).

Treatment and experimental design

The treatment tested pure nutmeg essential oil (NEO) and Petrogenol (800 g/l methyl eugenol) as an attractant and sterilized water as a control. The research was arranged to randomize block design with 10 replicates. Each two of the treatment units were put diagonally in the city area of Denpasar. Modified Steiner traps made from transparent plastic bottles (measuring 8 cm in diameter and 32 cm in length) (Fig. 1) were used to collect the fruit flies attracted on NEO and Petrogenol in the field (International Atomic Energy Agency, 2003).

Into trap 1 cotton was inserted containing 1.5 ml

sterilized water, while trap-2 cotton was inserted containing 1.5 ml NEO, and into trap-3 cotton was inserted containing 1.5 ml Petrogenol. Each trap was hung on the part of the plant about 2 to 3 meters above the soil surface. The distance of the inter-treatment trap (trap-1, trap-2, and trap-3) was about 40 m to 50 m, while the distance of inter replication traps was about three km. The observation of fruit flies-captured by the traps was conducted every day (24 hours after treatment).



Fig. 1. Types of fruit fly trap applied in this study

The parameters observed

The variables observed were the type of species and composition, number of fruit flies captured, and activity period of NEO and Petrogenol in the field. The analysis of the chemical composition of NEO was also done in this research using GC-MS.

Data analysis

The result of the parameters observed were statistically analyzed with SPSS program version 23.0 by ANOVA and comparing means by LSD test (Rem *et al.*, 2020).

Results and Discussion

Attractants are substances formulated to attract flies. In these studies, we used Petrogenol and NEO as attractants. They had a prominent effect on attracting flies, especially for the Tephritidae family. Average of fruit flies captured per day on the traps with Petrogenol and NEO as an attractant is 17.04 adults and 12.22 adults, respectively. These effects

are significant compared with control ($P < 0.05$) (Table 1). All of the adult fruit flies captured on the traps were male. There are three species of fruit flies identified that are *Bactrocera carambolae*, *B. dorsalis*, and *B. umbrosa* (Fig. 2). This phenomenon indicated that nutmeg essential oil can strongly attract fruit flies.

Based on *Gas Chromatography-Mass Spectrophotometer* (GC-MS) analysis indicated that one of the chemicals components of NEO was 8.33% methyl eugenol (Table 2). According to de Kogel *et al.* (2007), the chemical substance of methyl eugenol was able to attract around 17 species of fruit flies such as *B. araceae*, *B. carambolae*, *B. dorsalis*, *B. exornata*, *B. indonesiae*, *B. latilineola*, *B. muiri*, *B. nigrita*, *B. omlipitalis*, *B. papayae*, *B. platamus*, *B. raiensis*, *B. sulawesiae*, *B. thailandica*, *B. unimaculata*, and *B. verbascifoliae*. Vargas *et al.* (2000) stated that males of *B. dorsalis* were very attracted to methyl eugenol. According to Ghanim, (2013), many species of *Bactrocera* males after consumption of the methyl eugenol can enhance their mating competitiveness.

Table 1. Average of fruit flies captured per day on each attractant

Treatment	Average of fruit flies captured per day (adults)
Petrogenol	17.04 ^a
NEO	12.22 ^b
Distillated water	0.00 ^c

Note: NEO : *Nutmeg Essential Oil*. Different letter notations in the column indicated a significant difference between treatments ($P < 0.05$) based on the LSD test.

Figure 2 showed that species of *B. carambolae* and *B. dorsalis* were dominantly captured than *B. umbrosa*. This phenomenon was caused by both of the species having wider host plant range available in the field compare with *B. umbrosa*. According to Clarke *et al.* (2005); Maryati *et al.* (2008); Sauers-Muller, (2005) that *B. carambolae* has the widest host range and wide geographic distribution. As reported by Chinajariyawong *et al.* (2000), *B. carambolae* and *B. papayae* have the host plants more than *B. umbrosa*.

The activity period of NEO and Petrogenol attracting fruit flies in the field of more than 43 days is shown in Figure 3. The results showed that the number of fruit flies caught in traps with Petrogenol




Fig. 2. Type and composition of species fruit flies captured on the traps contain Petrogenol and NEO treatments

was higher than NEO. This phenomenon can be caused by the higher content of methyl eugenol in Petrogenol treatment 800 g/l when compared to NEO treatment which is only 8.33% (Table 2). The higher content of Methyl eugenol Petrogenol had a stronger effect on attracting male fruit flies than NEO.




Fig. 3. Activity period of attractant to attract fruit flies number per day in the field study

Conclusion

The nutmeg essential oil (*Myristica fragrans*) has the ability to attract only the male fruit flies (*Bactrocera*

dorsalis complex). There are three species of fruit flies identified that are *Bactrocera carambolae*, *B. papayae*, and *B. umbrosa*. Among the those species, *B. carambolae* and *B. dorsalis* were dominantly attracted. The results of GC-MS analysis showed that the compound 4-allyl-1,2-dimethoxybenzene-carboxylate (Methyl eugenol) was identified as a fruit fly attractant with a percentage of 8.33%. The activity period of nutmeg and petrogenol essential oil to attract fruit flies in the fields for more than 43 days shows that the number of fruit flies caught in the traps with petrogenol is higher than the essential oil of nutmeg.

The NEO material in this study has good development prospects in future research and still requires a lot of field trials so that it can reduce the use of synthetic pesticides.

Acknowledgement

The authors would like to of the Head Integrated Pest Management Laboratory (IPMLab), Faculty of

Table 2. Chemical composition of NEO analyzing with GC-MS

No.	Formula of molecule	Chemical components	Percentage (%)
1	C ₁₀ H ₁₆	Camphene	0.89
2	C ₁₀ H ₁₆	(+)-4-Carene	3.33
3	C ₁₀ H ₁₄	1-methyl-4-(1-methylethyl)benzene (Camphogen)	2.18
4	C ₁₀ H ₁₆	1-methyl-4-(1-methylethylidene)cyclohexene (Terpinolene)	2.91
5	C ₁₀ H ₁₀ O ₂	5-(1-propenyl)1,3-benzodioxole	1.13
6	C ₁₅ H ₂₄	á-cubebene	1.11
7	C ₁₁ H ₁₄ O ₂	4-allyl-1,2-dimethoxybenzene-carboxylate (Methyleugenol)	8.33
8	C ₁₅ H ₂₄	Caryophyllene	0.62
9	C ₁₁ H ₁₄ O ₂	1,2-dimethoxy-4-(1-propenyl)benzene (Methylisoeugenol)	5.02
10	C ₁₁ H ₁₂ O ₃	4-methoxy-6-(2-propenyl)1,3-benzodioxole (Myristicine)	6.58
11	C ₁₂ H ₁₆	1,2,3-trimethyl-5-(2-propenyl)benzene	3.0

Agriculture, Udayana University, Bali, Indonesia which has provided facilities and funding assistance.

References

- Afifah, N., 1, Kismiyati, Kenconoajati, H. 2020. Evaluation of aqueous extract of robusta coffee (*Coffea canephora*) leaves for controlling *Argulus japonicus* infestation on common carp seed. *IOP Conf. Series: Earth and Environmental Science*. 441 (2020) 012084. IOP Publishing. doi:10.1088/1755-1315/441/1/012084.
- Aktar, W., Sengupta, D. and Chowdhury, A. 2009. Impact of pesticides use in agriculture: their benefits and hazards. *Interdiscip Toxicol*. 2 : 1–12; doi: 10.2478/v10102-009-0001-7.
- Baser, K.H.C. and CRCbauer, G, eds. 2015. *Handbook of Essential Oils*. CRC Press.
- Bernardi, G., Kemmerich, M., Ribeiro, L.C., Adaime, M.B., Zanella, R. and Prestes, O.D. 2016. An effective method for pesticide residues determination in tobacco by GC-MS/MS and UHPLC-MS/MS employing acetonitrile extraction with low-temperature precipitation and d-SPE clean-up. *Talanta*. 161 : 40–47; doi:10.1016/j.talanta.2016.08.015.
- Butnariu, M. and Sarac, I. 2018. Essential Oils from Plants. J. Wan, ed *J Biotechnol Biomed Sci*. 1 : 35–43; doi:10.14302/issn.2576-6694.jbbs-18-2489.
- Chinajariyawong, A., Clarke, A.R., Jirasurat, M., Kritsaneepiboon, S., Lahey, H.A., Vijayasegaran, S. et al., 2000. Survey of opine parasitoids of fruit flies (Diptera: Tephritidae) in Thailand and Malaysia. *Raffles Bull Zool*. 48: 71–101.
- Clarke, A.R., Armstrong, K.F., Carmichael, A.E., Milne, J.R., Raghu, S. and Roderick, G.K. 2005. Invasive Phytophagous Pests Arising Through A Recent Tropical Evolutionary Radiation: The *Bactrocera dorsalis* Complex of Fruit Flies. *Annu Rev Entomol*. 50 : 293–319; doi: 10.1146/annurev.ento. 50.071803.130428.
- Damos, P., Colomar, L.A.E. and Ioriatti, C. 2015. Integrated fruit production and pest management in europe: The apple case study and how far we are from the original concept? *Insects*. 6 : 626–657; doi : 10.3390/insects6030626.
- de Kogel, W., Muryati, M. and Hasyim, A. 2007. Distribusi Spesies Lalat Buah di Sumatera Barat and Riau. *J Hortik*. 17 : 82161; doi:10.21082/jhort.v17n1.2007.p.
- Dhillon, M.K., Singh, R., Naresh, J.S. and Sharma, H.C. 2005. The melon fruit fly, *Bactrocera cucurbitae*: A review of its biology and management. *J Insect Sci*. 5; doi:10.1093/jis/5.1.40.
- Ghanim, M.N. 2013. Influence of Methyl Eugenol Diluted with Paraffin Oil on Male Annihilation Technique of Peach Fruit Fly, *Bactrocera zonata* (Saunders) (Diptera: Tephritidae). *Entomol Ornithol Herpetol Curr Res*. 02; doi:10.4172/2161-0983.1000114.
- Gupta, A.D., Bansal, V.K., Babu, V. and Maithil, N. 2013. Chemistry, antioxidant and antimicrobial potential of nutmeg (*Myristica fragrans* Houtt). *J Genet Eng Biotechnol*. 11 : 25–31; doi:10.1016/j.jgeb.2012.12.001.
- Ibrahim, I., Sudiana, I.K., Mukono, H.J. and Suhartono, Nugroho, H.S.W. 2020a. Awareness Program of Pesticides Used among Farmers using Difficulty-Usefulness Pyramid (A Suggestion for Health Laws and Policies Regarding the Use of Pesticides). *Indian Journal of Forensic Medicine & Toxicology*. 14 (3) : 1946–1951.
- Ibrahim, I., Sudiana, I.K., Mukono, H.J. and Suhartono, Nugroho, H.S.W. 2020b. Determination of Priority Elements of Vigilance in the Use of Pesticides based on Difficulty and Usefulness (A Supporting Study for Law and Policy in Health). *Indian Journal of Forensic Medicine & Toxicology*. 14(2) : 1615–1619.
- International Atomic Energy Agency. 2003. Area-Wide Fruit Fly Programmes Trapping Guidelines for Area-Wide. At Energy.
- Jaleel, W., Wang, D., Lei, Y., Qi, G., Chen, T. and Rizvi, S.A.H. 2020. Evaluating the repellent effect of four botanicals against two *Bactrocera* species on mangoes. *Peer J*. 8 : e8537; doi:10.7717/peerj.8537.
- Kardinan, A., Bintoro, M.H., Syakir, M. and Amin, A. 2020. Penggunaan Selasih Dalam Pengendalian Hama Lalat Buah Pada Mangga. *J Penelit Tanam Ind*. 15 : 101; doi:10.21082/jlitri.v15n3.2009.101-109.
- Kasman, K., Ishak, N.I., Hastutiek, P., Suprihati, E. and Mallongi, A. 2020. Identification of Active Compounds of Ethanol Extract of *Citrus amblycarpa* leaves by Analysis of Thin-layer Chromatography and Gas Chromatography-Mass Spectrometry as Bioinsecticide Candidates for Mosquitoes. *Open Access Macedonian Journal of Medical Sciences*. 2020 Sep 20 8(T2) : 1-6.https://doi.org/10.3889/oamjms.2020.5207
- Lanari, D., Marcotullio, M. and Neri, A. 2018. A Design of Experiment Approach for Ionic Liquid-Based Extraction of Toxic Components-Minimized Essential Oil from *Myristica fragrans* Houtt. *Fruits †. Molecules*. 23 : 2817; doi:10.3390/molecules23112817.
- Maryati, A., Hasyim, H. and Riska, R. 2008. Preferensi Spesies Lalat Buah Terhadap Atraktan Metil Eugenol Dan Cue-Lure Dan Populasinya Di Sumatera Barat Dan Riau. *J Hortik*. 18 : 85646; doi:10.21082/jhort.v18n2.2008.p.
- Muchtaridi, Subarnas, A., Apriyantono, A. and Mustarichie, R. 2010. Identification of Compounds in the Essential Oil of Nutmeg Seeds (*Myristica fragrans* Houtt.) That Inhibit Locomotor Activity in Mice. *Int J Mol Sci*. 11 : 4771–4781; doi:10.3390/ijms11114771.
- Ngowi, A.V.F., Mbise, T.J., Ijani, A.S.M., London, L. and

- Ajayi, O.C. 2007. Smallholder vegetable farmers in Northern Tanzania: Pesticides use practices, perceptions, cost and health effects. *Crop Prot.* 26 : 1617–1624; doi:10.1016/j.cropro.2007.01.008.
- Prastika, A.B., Sulmartiwi, L. and Lutfiyah, L. 2021. Effect of sublethal dose organophosphate pesticides on embryogenesis and hatching rate of silver rasbora eggs (*Rasbora argyrotaenia*). *IOP Conf. Series: Earth and Environmental Science.* 679 (2021) 012065 IOP Publishing. doi : 10.1088/1755-1315/679/1/012065
- Radonjic, S., Hrnèc, S. and Perovic, T. 2019. Overview of fruit flies important for fruit production on the Montenegro seacoast. *Biotechnol Agron Soc Environ.* 23 : 46–56; doi:10.25518/1780-4507.17776.
- Rem, P., Chiayvareesajja, S. and Suanyuk, N. 2020. Effects of temperature on growth performance and water quality in culture system of butter catfish (*Ompok bimaculatus*). *Songklanakarin J Sci Technol.* 42 : 1253–1258.
- Sanchez-Bayo, F. 2011. Impacts of Agricultural Pesticides on Terrestrial Ecosystems. In: *Ecological Impacts of Toxic Chemicals (Open Access)*. Bentham Science Publishers. 63–87.
- Sarwar, M., Hamed, M., Yousaf, M. and Hussain, M. 2014. Surveillance on Population Dynamics and Fruits Infestation of Tephritid Fruit Flies (Diptera: Tephritidae) in Mango (*Mangifera indica* L.) Orchards of Faisalabad, Pakistan. *Int J Sci Res Environ Sci.* 2 : 113–119; doi:10.12983/ijres-2014-p0113-0119.
- Setiawan, Koerniasari, Ngadino and Sudjarwo, S.A. 2017. Bioinsecticide Effect of *Pinus merkusii* Tree Bark Extract on *Aedes aegypti* larvae. *Journal of Young Pharmacists.* 9(1) : 127-130.doi:10.5530/jyp.2017.9.24.
- Sauers-Muller, A van. 2005. Host plants of the carambola fruit fly, *Bactrocera carambolae* Drew & Hancock (Diptera: Tephritidae), in Suriname, South America. *Neotrop Entomol.* 34 : 203–214; doi:10.1590/S1519-566X2005000200008.
- Singh, N.S., Sharma, R., Parween, T. and Patanjali, P.K. 2018. Pesticide Contamination and Human Health Risk Factor. In: *Modern Age Environmental Problems and their Remediation*. Springer International Publishing: Cham. 49–68.
- Siwi, S. and Hidayat, P. and Suputra. 2006. *Taksonomi dan Bioekologi Lalat Buah Penting di Indonesia (Diptera: Tephritidae)*. Balai Besar Penelitian dan Pengembangan Bioteknologi and Sumberdaya Genetik Pertanian: Bogor, Indonesia.
- Thanh Mai, N., Nga Thi Thanh, L., Jouni, H. and David, B. H. 2018. Pesticide use in vegetable production: A survey of Vietnamese farmers' knowledge. *Plant Prot Sci.* 54 : 203–214; doi:10.17221/69/2017-PPS.
- Vargas, R.I., Stark, J.D., Kido, M.H., Ketter, H.M. and Whitehand, L.C. 2000. Methyl Eugenol and Cue-Lure Traps for Suppression of Male Oriental Fruit Flies and Melon Flies (Diptera: Tephritidae) in Hawaii: Effects of Lure Mixtures and Weathering. *J Econ Entomol.* 93 : 81–87; doi:10.1603/0022-0493-93.1.81.
- White I., Elson-Harris, M. 1992. *Fruit flies of economic significance: their identification and bionomics*. CAB International Wallingford UK:International Institute of Entomology, London, SW7 5JR, UK.
- Wilson, C. and Tisdell, C. 2001. Why farmers continue to use pesticides despite environmental, health and sustainability costs. *Ecol Econ.* 39 : 449–462; doi:10.1016/S0921-8009(01)00238-5.